

**LIST OF TABLES**

Table 1.1 Dormant Season Agricultural Use of Diazinon by crop in Lower SJR Basin (1995-2002) in lbs. of a.i.

Table 1.2 Irrigation (in-season) Agricultural Use of Diazinon by Crop in Lower SJR (1995-2002) in lbs. of a.i.

Table 1.3 Dormant Season Agricultural Use of Chlorpyrifos by crop in Lower SJR Basin (1995-2002) in lbs. of a.i.

Table 1.4 Irrigation (in-season) Agricultural Use of Chlorpyrifos by Crop in Lower SJR (1995-2002) in lbs. of a.i.

Table 1.5 Annual Exceedances of Proposed Diazinon Acute Toxicity Target at the Mainstem Sites of the San Joaquin River (1991 – 2005)

Table 1.6 Annual Exceedances of Proposed Chlorpyrifos Acute Toxicity Water Quality Objective at the Mainstem Sites of the San Joaquin River (1991 – 2005)

Table 1.7 Basin Plan Method Analysis of Annual Exceedances of Combined Diazinon and Chlorpyrifos Toxicity at the Mainstem Sites of the San Joaquin River (1991 – 2005)

Table 1.8 Annual Exceedances of Proposed Diazinon Acute Toxicity Target at the Tributary Sites (1991 – 2005)

Table 1.9 Annual Exceedances of Proposed Chlorpyrifos Acute Toxicity Water Quality Objective at the Tributary Sites (1991 – 2005)

Table 1.10 Basin Plan Method Analysis of Annual Exceedances of Combined Diazinon and Chlorpyrifos Toxicity at the Tributary Sites (1991 – 2005)

Table 4.1 Water quality criteria for diazinon

Table 4.2 Water quality criteria for chlorpyrifos

Table 4.3 Summary of potential freshwater water quality objectives derived by alternate methods

Table 4.4 Comparison of historical data to the alternate water quality objectives

Table 4.5 Assessment of chlorpyrifos alternatives for their consistency with Porter-Cologne and other state and federal requirements

Table 4.6 Assessment of diazinon alternatives for their consistency with Porter-Cologne and other state and federal requirements

Table 4.7 River Reaches and Their Tributary Subareas

Table 4.8 Number and Magnitude of Observed Exceedances of Proposed Loading Capacity for Combined Diazinon and Chlorpyrifos Toxicity in the SJR (2000-2004)

Table 4.9 Number and Magnitude of Observed Exceedances of Proposed Loading Capacity for Diazinon in the SJR (2000-2004)

Table 4.10 Number and Magnitude of Observed Exceedances of Proposed Loading Capacity for Combined Diazinon and Chlorpyrifos Toxicity in SJR Tributaries (2000-2004)

Table 4.11 Number and Magnitude of Observed Exceedances of Proposed Loading Capacity for Diazinon in SJR Tributaries (2000-2004)

Table 5.1 Summary of Differences in Dormant Season Pest Management Costs

Table 5.2 Surface Irrigation - Initial Capital Cost and Recurring Maintenance Costs (from Burt et al. 2000)

Table 5.3 Sprinkler Irrigation - Initial Capital Cost and Recurring Maintenance Costs (from Burt et al. (2000)

Table 5.4 Micro-irrigation - Initial Capital and Maintenance Costs (from Burt et al (2000)

Table 5.5 Estimated Cost to Convert from Flood to Sprinkler Irrigation

Table 5.6 Drainage Practices Capital and Maintenance Costs

Table 5.7 Summary of Differences in Irrigation Season Pest Management Costs

Table 7.1 Summary of Public Workshops

**Table 1.1. Dormant Season Agricultural Use of Diazinon by crop in Lower SJR Basin (1995-2002) in lbs. of a.i.**

| Commodity | 1995   | 1996   | 1997   | 1998   | 1999   | 2000   | 2001   | 2002   | Average | % Average* |
|-----------|--------|--------|--------|--------|--------|--------|--------|--------|---------|------------|
| ALMOND    | 28,893 | 35,134 | 18,743 | 33,640 | 37,948 | 10,668 | 18,719 | 17,680 | 25,178  | 65.32%     |
| PEACH     | 7,383  | 6,518  | 4,599  | 5,353  | 5,552  | 4,022  | 4,068  | 5,499  | 5,374   | 13.94%     |
| APRICOT   | 6,622  | 3,945  | 920    | 2,712  | 2,350  | 2,516  | 113    | 113    | 2,411   | 6.26%      |
| PRUNE     | 2,676  | 1,269  | 1,213  | 486    | 1,851  | 1,273  | 821    | 2,840  | 1,554   | 4.03%      |
| APPLE     | 3,113  | 2,593  | 2,514  | 1,008  | 752    | 686    | 446    | 395    | 1,438   | 3.73%      |
| NECTARINE | 1,452  | 1,219  | 1,046  | 1,213  | 1,306  | 1,213  | 1,151  | 794    | 1,174   | 3.05%      |
| PLUM      | 1,259  | 953    | 786    | 779    | 681    | 837    | 982    | 456    | 842     | 2.18%      |
| TOTAL     | 51,398 | 51,631 | 29,821 | 45,191 | 50,440 | 21,215 | 26,300 | 27,777 | NA      | 98.51%     |

\* % Average values do not sum to 100% as crops with less than 1% average use are not shown

139

**Table 1.2. Irrigation (in-season) Agricultural Use of Diazinon by Crop in Lower SJR (1995-2002) in lbs. of a.i.**

| Commodity   | 1995   | 1996   | 1997   | 1998   | 1999   | 2000   | 2001   | 2002  | Average | % Average* |
|-------------|--------|--------|--------|--------|--------|--------|--------|-------|---------|------------|
| ALMOND      | 35,371 | 13,050 | 2,134  | 227    | 683    | 168    | 90     | 2     | 6,466   | 26.84%     |
| CANTALOUPE  | 2,963  | 3,185  | 4,297  | 877    | 2,977  | 2,163  | 2,797  | 2,653 | 2,739   | 11.37%     |
| PEACH       | 3,954  | 3,807  | 2,433  | 993    | 1,670  | 2,375  | 2,376  | 597   | 2,276   | 9.45%      |
| TOMATO      | 2,207  | 1,701  | 363    | 835    | 812    | 3,765  | 2,977  | 695   | 1,670   | 6.93%      |
| MELON       | 2,111  | 1,630  | 1,897  | 1,616  | 1,982  | 1,007  | 964    | 1,979 | 1,648   | 6.84%      |
| PRUNE       | 984    | 1,210  | 518    | 4,205  | 1,979  | 2,302  | 414    | 1,311 | 1,615   | 6.71%      |
| WALNUT      | 2,137  | 1,634  | 2,606  | 975    | 311    | 1,357  | 1,398  | 61    | 1,310   | 5.44%      |
| APRICOT     | 2,075  | 1,631  | 894    | 1,186  | 1,544  | 743    | 212    | 83    | 1,046   | 4.34%      |
| ALFALFA     | 3,099  | 3,456  | 177    | 307    | 1      | 0      | 0      | 0     | 880     | 3.65%      |
| APPLE       | 1,742  | 1,877  | 528    | 283    | 771    | 587    | 292    | 723   | 850     | 3.53%      |
| NECTARINE   | 1,451  | 1,140  | 569    | 430    | 727    | 1,282  | 750    | 113   | 808     | 3.35%      |
| PLUM        | 1,433  | 976    | 364    | 157    | 350    | 225    | 274    | 21    | 475     | 1.97%      |
| BEANS       | 498    | 538    | 845    | 254    | 10     | 829    | 100    | 0     | 384     | 1.59%      |
| WATERMELON  | 158    | 212    | 798    | 300    | 377    | 131    | 186    | 131   | 287     | 1.19%      |
| GRAPE, WINE | 621    | 281    | 268    | 82     | 202    | 40     | 68     | 381   | 243     | 1.01%      |
| TOTAL       | 60,804 | 36,328 | 18,691 | 12,727 | 14,396 | 16,974 | 12,898 | 8,750 | NA      | 94.21%     |

\* % Average values do not sum to 100% as crops with less than 1% average use are not shown

**Table 1.3. Dormant Season Agricultural Use of Chlorpyrifos by crop in Lower SJR Basin (1995-2002) in lbs. of a.i.**

| Commodity | 1995   | 1996   | 1997  | 1998   | 1999   | 2000  | 2001   | 2002   | Average | % Average* |
|-----------|--------|--------|-------|--------|--------|-------|--------|--------|---------|------------|
| ALMOND    | 9,668  | 10,430 | 3,966 | 6,625  | 8,109  | 1,520 | 7,509  | 7,844  | 6,959   | 52.65%     |
| APPLE     | 4,713  | 3,006  | 2,867 | 2,626  | 2,433  | 1,751 | 1,415  | 1,190  | 2,500   | 18.92%     |
| PEACH     | 2,754  | 1,803  | 1,066 | 785    | 1,040  | 832   | 2,120  | 3,002  | 1,675   | 12.68%     |
| ALFALFA   | 1,868  | 427    | 816   | 15     | 70     | 2,266 | 136    | 105    | 713     | 5.39%      |
| FIG       | 0      | 0      | 0     | 0      | 259    | 0     | 4,871  | 0      | 641     | 4.85%      |
| NECTARINE | 48     | 60     | 319   | 241    | 407    | 244   | 97     | 32     | 181     | 1.37%      |
| GRAPE     | 0      | 0      | 704   | 40     | 0      | 203   | 214    | 24     | 148     | 1.12%      |
| TOTAL     | 19,051 | 15,726 | 9,738 | 10,332 | 12,318 | 6,816 | 16,362 | 12,197 | NA      | 96.98%     |

\* % Average values do not sum to 100% as crops with less than 1% average use are not shown

**Table 1.4. Irrigation (in-season) Agricultural Use of Chlorpyrifos by Crop in Lower SJR (1995-2002) in lbs. of a.i.**

| Commodity    | 1995    | 1996    | 1997    | 1998    | 1999    | 2000    | 2001    | 2002    | Average | % Average* |
|--------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|------------|
| ALMOND       | 71,339  | 93,617  | 104,911 | 109,162 | 76,902  | 88,371  | 76,374  | 55,776  | 84,556  | 39.23%     |
| COTTON       | 116,733 | 24,561  | 44,867  | 23,104  | 18,960  | 17,656  | 20,716  | 5,666   | 34,033  | 15.79%     |
| ALFALFA      | 59,720  | 46,583  | 36,515  | 40,857  | 22,684  | 25,180  | 17,163  | 14,682  | 32,923  | 15.28%     |
| WALNUT       | 34,281  | 34,829  | 31,196  | 28,923  | 26,436  | 24,160  | 29,588  | 26,002  | 29,427  | 13.65%     |
| CORN         | 13,250  | 7,403   | 11,551  | 8,812   | 13,110  | 12,932  | 7,475   | 7,077   | 10,201  | 4.73%      |
| APPLE        | 10,710  | 9,334   | 9,955   | 12,542  | 4,459   | 2,290   | 662     | 66      | 6,252   | 2.90%      |
| SUGARBEET    | 3,455   | 3,478   | 4,842   | 6,505   | 7,216   | 3,234   | 3,152   | 2,327   | 4,276   | 1.98%      |
| ORANGE       | 4,060   | 2,937   | 1,782   | 5,092   | 7,010   | 2,059   | 2,936   | 3,885   | 3,720   | 1.73%      |
| SWEET POTATO | 1,122   | 1,794   | 2,691   | 3,061   | 5,571   | 3,964   | 5,539   | 721     | 3,058   | 1.42%      |
| GRAPE        | 0       | 514     | 1,117   | 5,964   | 3,808   | 2,243   | 5,253   | 2,569   | 2,684   | 1.25%      |
| TOTAL        | 314,670 | 225,050 | 249,427 | 244,022 | 186,156 | 183,089 | 168,858 | 118,771 | NA      | 97.96%     |

\* % Average values do not sum to 100% as crops with less than 1% average use are not shown.

**Table 1.5. Annual Exceedances of Proposed Diazinon Acute Toxicity Target at the Mainstem Sites of the San Joaquin River (1991 – 2005)**

| Site Name                         | 1991                                       | 1992               | 1993              | 1994              | 1995            | 1996 | 1997            | 1998            | 1999            | 2000            | 2001              | 2002            | 2003            | 2004            | 2005            |
|-----------------------------------|--|--------------------|-------------------|-------------------|-----------------|------|-----------------|-----------------|-----------------|-----------------|-------------------|-----------------|-----------------|-----------------|-----------------|
| SJR near Vernalis                 | <b>0%</b> <sup>a</sup><br>160 <sup>b</sup> | <b>2.5%</b><br>200 | <b>12%</b><br>266 | <b>16%</b><br>120 | <b>0%</b><br>14 | NS   | <b>0%</b><br>34 | <b>0%</b><br>43 | <b>0%</b><br>44 | <b>0%</b><br>74 | <b>14%</b><br>65  | <b>0%</b><br>11 | <b>0%</b><br>36 | <b>0%</b><br>31 | <b>0%</b><br>38 |
| SJR at Maze Blvd.                 | <b>0%</b><br>3                             | <b>40%</b><br>5    | <b>50%</b><br>2   | NS                | NS              | NS   | NS              | NS              | NS              | NS              | <b>0%</b><br>20   | NS              | NS              | NS              | NS              |
| SJR at Crows Landing              | NS   | NS                 | NS                | NS                | NS              | NS   | NS              | NS              | NS              | NS              | <b>0%</b><br>23   | <b>0%</b><br>6  | <b>0%</b><br>34 | <b>0%</b><br>15 | NS              |
| SJR near Patterson                | NS   | NS                 | NS                | <b>0%</b><br>5    | NS              | NS   | NS              | NS              | NS              | <b>0%</b><br>1  | <b>0%</b><br>40   | <b>0%</b><br>9  | NS              | NS              | <b>0%</b><br>8  |
| SJR near Newman                   | <b>0%</b><br>5                             | <b>4.5%</b><br>22  | <b>50%</b><br>2   | NS                | NS              | NS   | NS              | NS              | NS              | NS              | NS                | NS              | NS              | NS              | NS              |
| SJR near Stevinson at Lander Ave. | <b>0%</b><br>3                             | <b>0%</b><br>5     | <b>50%</b><br>2   | <b>0%</b><br>1    | NS              | NS   | NS              | NS              | NS              | <b>5%</b><br>20 | <b>2.2%</b><br>43 | NS              | <b>0%</b><br>16 | NS              | <b>0%</b><br>7  |

Proposed Diazinon Acute Toxicity Target = 0.16 µg/L

<sup>a</sup>Percent of samples for the year that exceed the proposed diazinon acute toxicity target value.

<sup>b</sup>Total number of samples analyzed for diazinon during the year.

NS = No samples analyzed during the year.

**Table 1.6. Annual Exceedances of Proposed Chlorpyrifos Acute Toxicity Water Quality Objective at the Mainstem Sites of the San Joaquin River (1991 – 2005)**

| Site Name                         | 1991                                       | 1992                | 1993               | 1994               | 1995              | 1996 | 1997             | 1998           | 1999            | 2000               | 2001              | 2002             | 2003              | 2004              | 2005            |
|-----------------------------------|--|---------------------|--------------------|--------------------|-------------------|------|------------------|----------------|-----------------|--------------------|-------------------|------------------|-------------------|-------------------|-----------------|
| SJR near Vernalis                 | <b>0%</b> <sup>a</sup><br>174 <sup>b</sup> | <b>0.50%</b><br>179 | <b>4.5%</b><br>155 | <b>1.9%</b><br>102 | <b>7.1%</b><br>14 | NS   | <b>0%</b><br>320 | <b>0%</b><br>9 | <b>0%</b><br>43 | <b>2.4%</b><br>333 | <b>0%</b><br>542  | <b>0%</b><br>145 | <b>0%</b><br>39   | <b>3.2%</b><br>31 | <b>0%</b><br>23 |
| SJR at Maze Blvd.                 | <b>0%</b><br>3                             | <b>0%</b><br>5      | <b>0%</b><br>2     | NS                 | NS                | NS   | NS               | NS             | NS              | NS                 | <b>0%</b><br>20   | NS               | NS                | NS                | NS              |
| SJR at Crows Landing              | NS   | NS                  | NS                 | NS                 | NS                | NS   | NS               | NS             | NS              | NS                 | <b>8.7%</b><br>23 | <b>50%</b><br>6  | <b>3%</b><br>33   | <b>6.7%</b><br>15 | NS              |
| SJR near Patterson                | NS   | NS                  | NS                 | <b>20%</b><br>5    | NS                | NS   | NS               | NS             | NS              | <b>0%</b><br>1     | <b>0%</b><br>40   | <b>12%</b><br>8  | NS                | NS                | <b>0%</b><br>9  |
| SJR near Newman                   | <b>0%</b><br>28                            | <b>0%</b><br>28     | <b>0%</b><br>2     | NS                 | NS                | NS   | NS               | NS             | NS              | NS                 | NS                | NS               | NS                | NS                | NS              |
| SJR near Stevinson at Lander Ave. | <b>0%</b><br>3                             | <b>0%</b><br>5      | <b>50%</b><br>2    | <b>0%</b><br>1     | NS                | NS   | NS               | NS             | NS              | <b>5%</b><br>20    | <b>0%</b><br>43   | NS               | <b>5.9%</b><br>17 | NS                | <b>0%</b><br>7  |

Proposed Chlorpyrifos Acute Toxicity Water Quality Objective = 0.025 µg/L

<sup>a</sup>Percent of samples for the year that equal or exceed the proposed chlorpyrifos acute toxicity water quality objective value.<sup>b</sup>Total number of samples analyzed for chlorpyrifos during the year.

NS = No samples analyzed during the year.

**Table 1.7. Basin Plan Method Analysis of Annual Exceedances of Combined Diazinon and Chlorpyrifos Toxicity at the Mainstem Sites of the San Joaquin River (1991 – 2005)**

| Site Name                         | 1991                                       | 1992              | 1993              | 1994              | 1995              | 1996           | 1997            | 1998              | 1999              | 2000              | 2001              | 2002            | 2003              | 2004              | 2005              |
|-----------------------------------|--|-------------------|-------------------|-------------------|-------------------|----------------|-----------------|-------------------|-------------------|-------------------|-------------------|-----------------|-------------------|-------------------|-------------------|
| SJR near Vernalis                 | <b>0%</b> <sup>a</sup><br>169 <sup>b</sup> | <b>3%</b><br>200  | <b>15%</b><br>264 | <b>22%</b><br>103 | <b>7.1%</b><br>14 | <b>0%</b><br>3 | <b>0%</b><br>35 | <b>2.4%</b><br>42 | <b>2.4%</b><br>42 | <b>1.4%</b><br>71 | <b>19%</b><br>64  | <b>0%</b><br>12 | <b>0%</b><br>42   | <b>2.7%</b><br>36 | <b>4.2%</b><br>24 |
| SJR at Maze Blvd.                 | <b>0%</b><br>2                             | <b>20%</b><br>5   | <b>50%</b><br>2   | NS                | NS                | NS             | NS              | NS                | NS                | NS                | <b>0%</b><br>20   | NS              | NS                | NS                | NS                |
| SJR at Crows Landing              | NS   | NS                | NS                | NS                | NS                | NS             | NS              | NS                | NS                | NS                | <b>8.7%</b><br>23 | <b>38%</b><br>8 | <b>2.8%</b><br>36 | <b>4.0%</b><br>25 | NS                |
| SJR near Patterson                | NS   | NS                | NS                | <b>20%</b><br>5   | NS                | NS             | NS              | NS                | NS                | <b>0%</b><br>1    | <b>0%</b><br>40   | <b>11%</b><br>9 | NS                | NS                | <b>0%</b><br>9    |
| SJR near Newman                   | <b>4.5%</b><br>22                          | <b>4.5%</b><br>22 | <b>50%</b><br>2   | NS                | NS                | NS             | NS              | NS                | NS                | NS                | NS                | NS              | NS                | NS                | NS                |
| SJR near Stevinson at Lander Ave. | <b>0%</b><br>3                             | <b>0%</b><br>5    | <b>50%</b><br>2   | <b>0%</b><br>1    | NS                | NS             | NS              | NS                | NS                | <b>5%</b><br>20   | <b>14%</b><br>43  | NS              | <b>5.9%</b><br>17 | NS                | NS                |

<sup>a</sup>Percent of samples for the year for which the combined (additive) toxicity value equals or exceeds 1.0.

<sup>b</sup>Total number of samples analyzed for chlorpyrifos and/or diazinon during the year.

NS = No samples analyzed during the year. Acute diazinon WQO = 0.16µg/L; acute chlorpyrifos WQO = 0.025 µg/L.

**Table 1.8 Annual Exceedances of Proposed Diazinon Acute Toxicity Target at the Tributary Sites of the San Joaquin River**

| Site Name                              | 1991                                     | 1992              | 1993              | 1994             | 1995              | 1996               | 1997              | 1998            | 1999            | 2000             | 2001               | 2002             | 2003            | 2004              | 2005             |
|--|--|-------------------|-------------------|------------------|-------------------|--------------------|-------------------|-----------------|-----------------|------------------|--------------------|------------------|-----------------|-------------------|------------------|
| Stanislaus River at Caswell State Park | <b>0%</b> <sup>a</sup><br>6 <sup>b</sup> | <b>0%</b><br>13   | <b>0%</b><br>2    | <b>0%</b><br>1   | NS                | NS                 | NS                | NS              | NS              | <b>0%</b><br>20  | <b>0%</b><br>41    | <b>0%</b><br>9   | <b>0%</b><br>31 | <b>3.2%</b><br>31 | <b>17%</b><br>23 |
| Tuolumne River at Shiloh Road          | <b>0%</b><br>2                           | <b>13%</b><br>15  | <b>50%</b><br>2   | <b>100%</b><br>1 | NS                | NS                 | NS                | NS              | NS              | <b>0%</b><br>20  | <b>2.01%</b><br>49 | <b>0%</b><br>20  | <b>0%</b><br>32 | <b>0%</b><br>31   | <b>13%</b><br>22 |
| Del Puerto Creek at Vineyard Road      | <b>9%</b><br>11                          | <b>27%</b><br>18  | <b>0%</b><br>2    | <b>0%</b><br>1   | NS                | NS                 | NS                | NS              | NS              | <b>45%</b><br>11 | <b>0%</b><br>23    | <b>0%</b><br>16  | <b>0%</b><br>35 | NS                | NS               |
| Orestimba Creek at River Road          | <b>12%</b><br>8                          | <b>21%</b><br>66  | <b>18%</b><br>50  | <b>0%</b><br>1   | <b>0%</b><br>1    | <b>2.8%</b><br>254 | <b>13%</b><br>160 | <b>0%</b><br>32 | <b>0%</b><br>36 | <b>11%</b><br>46 | <b>0%</b><br>45    | <b>18%</b><br>27 | <b>0%</b><br>33 | NS                | NS               |
| Merced River at River Road             | <b>0%</b><br>4                           | <b>6.2%</b><br>16 | <b>9.5%</b><br>42 | <b>14%</b><br>50 | <b>3.6%</b><br>28 | NS                 | <b>0%</b><br>10   | <b>0%</b><br>15 | <b>0%</b><br>15 | <b>0%</b><br>32  | <b>6.7%</b><br>45  | <b>0%</b><br>8   | <b>0%</b><br>32 | <b>0%</b><br>23   | <b>0%</b><br>14  |
| Mud Slough near Gustine                | <b>0%</b><br>3                           | <b>0%</b><br>5    | <b>50%</b><br>2   | <b>0%</b><br>1   | NS                | NS                 | NS                | NS              | <b>0%</b><br>1  | NS               | <b>4.5%</b><br>22  | NS               | NS              | NS                | NS               |
| Salt Slough at Lander Avenue           | <b>0%</b><br>4                           | <b>19%</b><br>16  | <b>11%</b><br>28  | <b>0%</b><br>1   | NS                | NS                 | NS                | NS              | NS              | NS               | <b>4.5%</b><br>22  | NS               | <b>0%</b><br>17 | NS                | NS               |
| TID Lateral 5 (Harding Drain)          | <b>0%</b><br>7                           | <b>17%</b><br>41  | <b>60%</b><br>5   | <b>0%</b><br>1   | NS                | NS                 | NS                | NS              | NS              | <b>0%</b><br>11  | <b>0%</b><br>2     | NS               | NS              | NS                | NS               |
| Ingram/Hospital Creeks at River Road   | <b>25%</b><br>12                         | <b>32%</b><br>19  | <b>100%</b><br>2  | <b>0%</b><br>2   | NS                | NS                 | NS                | NS              | NS              | NS               | <b>25%</b><br>4    | NS               | NS              | NS                | NS               |
| Spanish Grant Drain near Patterson     | <b>0%</b><br>6                           | <b>7.1%</b><br>14 | <b>50%</b><br>2   | <b>0%</b><br>1   | NS                | NS                 | NS                | NS              | NS              | NS               | <b>0%</b><br>2     | NS               | NS              | NS                | NS               |

Proposed Diazinon Acute Toxicity Target = 0.16 µg/L

<sup>a</sup>Percent of samples for the year that exceed the proposed diazinon acute toxicity target value.<sup>b</sup>Total number of samples analyzed for diazinon during the year.

NS = No samples analyzed during the year.



**Table 1.9 Annual Exceedances of Proposed Chlorpyrifos Acute Toxicity Water Quality Objective at the Tributary Sites (1991 – 2005)**

| Site Name                              | 1991             | 1992              | 1993             | 1994             | 1995             | 1996              | 1997              | 1998             | 1999              | 2000              | 2001              | 2002              | 2003              | 2004              | 2005             |
|--|------------------|-------------------|------------------|------------------|------------------|-------------------|-------------------|------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|
| Stanislaus River at Caswell State Park | <b>0%</b><br>6   | <b>0%</b><br>13   | <b>0%</b><br>2   | <b>0%</b><br>1   | NS               | NS                | NS                | NS               | NS                | <b>0%</b><br>20   | <b>4.9%</b><br>41 | <b>0%</b><br>9    | <b>6.1%</b><br>33 | <b>3.1%</b><br>32 | <b>0%</b><br>23  |
| Tuolumne River at Shiloh Road          | <b>0%</b><br>2   | <b>6.7%</b><br>15 | <b>0%</b><br>2   | <b>0%</b><br>1   | NS               | NS                | NS                | NS               | NS                | <b>0%</b><br>20   | <b>0%</b><br>49   | <b>4.8%</b><br>21 | <b>3.0%</b><br>33 | <b>3.2%</b><br>31 | <b>14%</b><br>14 |
| Del Puerto Creek at Vineyard Road      | <b>22%</b><br>9  | <b>15%</b><br>20  | <b>0%</b><br>4   | <b>100%</b><br>1 | NS               | NS                | NS                | NS               | NS                | <b>0%</b><br>11   | <b>22%</b><br>23  | <b>19%</b><br>16  | <b>14%</b><br>35  | NS                | NS               |
| Orestimba Creek at River Road          | <b>50%</b><br>8  | <b>28%</b><br>65  | <b>27%</b><br>51 | <b>100%</b><br>1 | <b>100%</b><br>1 | <b>32%</b><br>243 | <b>29%</b><br>198 | <b>12%</b><br>32 | <b>5.4%</b><br>37 | <b>2.1%</b><br>47 | <b>18%</b><br>45  | <b>7.4%</b><br>27 | <b>16%</b><br>32  | NS                | NS               |
| Merced River at River Road             | <b>0%</b><br>4   | <b>19%</b><br>16  | <b>46%</b><br>41 | <b>26%</b><br>46 | <b>0%</b><br>33  | NS                | <b>0%</b><br>10   | <b>0%</b><br>15  | <b>0%</b><br>15   | <b>0%</b><br>31   | <b>2.3%</b><br>43 | <b>0%</b><br>9    | <b>0%</b><br>32   | <b>0%</b><br>24   | <b>33%</b><br>15 |
| TID Lateral 5 (Harding Drain)          | <b>57%</b><br>7  | <b>32%</b><br>40  | <b>28%</b><br>7  | <b>0%</b><br>1   | NS               | NS                | NS                | NS               | NS                | <b>0%</b><br>9    | <b>0%</b><br>2    | NS                | NS                | NS                | NS               |
| Ingram/Hospital Creeks at River Road   | <b>27%</b><br>11 | <b>9.5%</b><br>21 | <b>0%</b><br>2   | <b>0%</b><br>2   | NS               | NS                | NS                | NS               | NS                | <b>0%</b><br>4    | NS                | NS                | NS                | NS                | NS               |
| Spanish Grant Drain near Patterson     | <b>83%</b><br>6  | <b>57%</b><br>14  | <b>0%</b><br>3   | <b>25%</b><br>4  | NS               | NS                | NS                | NS               | NS                | NS                | <b>0%</b><br>2    | NS                | NS                | NS                | NS               |
| Mud Slough near Gustine                | <b>0%</b><br>3   | <b>0%</b><br>5    | <b>0%</b><br>2   | <b>0%</b><br>1   | NS               | NS                | NS                | NS               | <b>0%</b><br>1    | NS                | <b>4.5%</b><br>22 | NS                | NS                | NS                | NS               |
| Salt Slough at Lander Ave.             | <b>0%</b><br>3   | <b>6.7%</b><br>15 | <b>21%</b><br>28 | <b>0%</b><br>1   | NS               | NS                | NS                | NS               | NS                | NS                | <b>14%</b><br>22  | <b>100%</b><br>1  | <b>0%</b><br>18   | NS                | NS               |

Proposed Chlorpyrifos Acute Toxicity Water Quality Objective = 0.025 µg/L

<sup>a</sup>Percent of samples for the year that equal or exceed the proposed chlorpyrifos acute toxicity water quality objective value.<sup>b</sup>Total number of samples analyzed for chlorpyrifos during the year.

NS = No samples analyzed during the year.

**Table 1.10. Basin Plan Method Analysis of Annual Exceedances of Combined Diazinon and Chlorpyrifos Toxicity at the Tributary Sites (1991 – 2005)**

| Site Name                              | 1991                                     | 1992             | 1993             | 1994             | 1995              | 1996              | 1997              | 1998             | 1999              | 2000              | 2001              | 2002              | 2003              | 2004              | 2005             |
|--|--|------------------|------------------|------------------|-------------------|-------------------|-------------------|------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|
| Stanislaus River at Caswell State Park | <b>0%</b> <sup>a</sup><br>5 <sup>b</sup> | <b>0%</b><br>13  | <b>0%</b><br>2   | <b>0%</b><br>1   | NS                | NS                | NS                | NS               | NS                | <b>0%</b><br>20   | <b>4.1%</b><br>41 | <b>0%</b><br>9    | <b>5.7%</b><br>34 | <b>13%</b><br>30  | <b>30%</b><br>23 |
| Tuolumne River at Shiloh Road          | <b>0%</b><br>2                           | <b>13%</b><br>15 | <b>50%</b><br>2  | <b>100%</b><br>1 | NS                | NS                | NS                | NS               | NS                | <b>4.8%</b><br>21 | <b>10%</b><br>49  | <b>4.5%</b><br>22 | <b>0%</b><br>36   | <b>6.4%</b><br>31 | <b>17%</b><br>23 |
| Del Puerto Creek at Vineyard Road      | <b>40%</b><br>10                         | <b>50%</b><br>18 | <b>0%</b><br>2   | <b>100%</b><br>1 | NS                | NS                | NS                | NS               | NS                | <b>45%</b><br>11  | <b>22%</b><br>23  | <b>17%</b><br>16  | <b>14%</b><br>35  | NS                | NS               |
| Orestimba Creek at River Road          | <b>50%</b><br>8                          | <b>58%</b><br>66 | <b>40%</b><br>50 | <b>100%</b><br>1 | <b>100%</b><br>1  | <b>40%</b><br>244 | <b>46%</b><br>132 | <b>12%</b><br>32 | <b>5.6%</b><br>36 | <b>15%</b><br>46  | <b>22%</b><br>46  | <b>22%</b><br>31  | <b>14%</b><br>35  | NS                | NS               |
| Merced River at River Road             | <b>0%</b><br>4                           | <b>19%</b><br>16 | <b>48%</b><br>42 | <b>30%</b><br>50 | <b>3.4%</b><br>29 | NS                | <b>0%</b><br>10   | <b>0%</b><br>15  | <b>0%</b><br>15   | <b>0%</b><br>32   | <b>9.1%</b><br>44 | <b>0%</b><br>9    | <b>0%</b><br>33   | <b>0%</b><br>26   | <b>21%</b><br>14 |
| Mud Slough near Gustine                | <b>0%</b><br>3                           | <b>0%</b><br>5   | <b>50%</b><br>2  | <b>0%</b><br>1   | NS                | NS                | NS                | NS               | <b>0%</b><br>1    | NS                | <b>9%</b><br>22   | NS                | NS                | NS                | NS               |
| Salt Slough at Lander Ave.             | <b>0%</b><br>3                           | <b>25%</b><br>16 | <b>36%</b><br>28 | <b>0%</b><br>1   | NS                | NS                | NS                | NS               | NS                | NS                | <b>18%</b><br>22  | <b>0%</b><br>1    | <b>0%</b><br>18   | NS                | NS               |
| TID Lateral 5 (Harding Drain)          | <b>57%</b><br>7                          | <b>46%</b><br>41 | <b>80%</b><br>5  | <b>0%</b><br>1   | NS                | NS                | NS                | NS               | <b>0%</b><br>1    | <b>0%</b><br>9    | <b>0%</b><br>2    | NS                | NS                | NS                | NS               |
| Ingram/Hospital Creeks at River Road   | <b>54%</b><br>11                         | <b>37%</b><br>19 | <b>100%</b><br>2 | <b>0%</b><br>2   | NS                | NS                | NS                | NS               | NS                | NS                | <b>25%</b><br>4   | NS                | NS                | NS                | NS               |
| Spanish Grant Drain near Patterson     | <b>83%</b><br>6                          | <b>57%</b><br>14 | <b>50%</b><br>2  | <b>100%</b><br>1 | NS                | NS                | NS                | NS               | NS                | NS                | <b>0%</b><br>2    | NS                | NS                | NS                | NS               |

<sup>a</sup>Percent of samples for the year for which the combined (additive) toxicity value equals or exceeds 1.0.<sup>b</sup>Total number of samples analyzed for chlorpyrifos and/or diazinon during the year.

NS = No samples analyzed during the year.

**Table 4.1. Water quality criteria for diazinon**

| <b>Aquatic Life Criteria for Surface Water</b>  | <b>µg/L</b> |
|---|-------------|
| CDFG Aquatic Life Criteria for freshwater – 4 day average concentration                               | 0.05        |
| CDFG Aquatic Life Criteria for freshwater – 1 hour maximum concentration                              | 0.08        |
| Recalculated CDFG Aquatic Life Criteria for freshwater – 4 day average concentration                  | 0.10        |
| Recalculated CDFG Aquatic Life Criteria for freshwater – 1 hour maximum concentration                 | 0.16        |
| EPA Draft Aquatic Life Criteria for freshwater – 4 day average concentration                          | 0.10        |
| EPA Draft Aquatic Life Criteria for freshwater – 1 hour maximum concentration                         | 0.10        |
| Australian and New Zealand trigger values (95% protection- based on NOEC)                             | 0.010       |
| Australian and New Zealand trigger values (99% protection – based on NOEC)                            | 0.00003     |
| 1/10 <sup>th</sup> Species mean average value ( <i>Ceriodaphnia dubia</i> ) <sup>1</sup> (Basin Plan) | 0.044       |
| <b>Human Health Criteria for Drinking Water</b>   |             |
| US EPA Suggested No Adverse Response Levels (SNARL) for non-cancer toxicity                           | 0.600       |
| California Department of Health Services State Action Level for Toxicity                              | 6.000       |
| National Academy of Sciences SNARL for non-cancer toxicity  | 14.000      |
| Canadian Environmental Quality Guidelines   | 20.000      |

**Table 4.2 Water quality criteria for chlorpyrifos**

| <b>Aquatic Life Criteria for Surface Water</b>  | <b>µg/L</b> |
|---|-------------|
| CDFG Aquatic Life Criteria for freshwater – 4 day average concentration                               | 0.014       |
| CDFG Aquatic Life Criteria for freshwater – 1 hour maximum concentration                              | 0.02        |
| EPA Draft Aquatic Life Criteria for freshwater – 4 day average concentration                          | 0.041       |
| EPA Draft Aquatic Life Criteria for freshwater – 1 hour maximum concentration                         | 0.083       |
| Canadian Environmental Quality Guidelines   | 0.0035      |
| Australian and New Zealand trigger values (95% protection based on NOEC)                              | 0.010       |
| Australian and New Zealand trigger values (99% protection based on NOEC)                              | 0.00004     |
| 1/10 <sup>th</sup> Species mean average value ( <i>Ceriodaphnia dubia</i> ) <sup>2</sup> (Basin Plan) | 0.006       |
| <b>Human Health Criteria for Drinking Water</b>   |             |
| US EPA Suggested No Adverse Response Levels (SNARL) for non-cancer toxicity                           | 20.000      |
| Canadian Environmental Quality Guidelines   | 90.000      |
| <b>Agriculture-Livestock</b>  |             |
| Canadian Environmental Quality Guidelines   | 24.000      |

Sources: Marshack, 2003; Canadian Council of Ministers of the Environment, 2002; US EPA 2003; Siepmann and Finlayson, 2000; Finlayson, 2004; Australian and New Zealand Environment and Conservation Council and Agriculture and Resource Management Council of Australia and New Zealand, 2000.

<sup>1</sup> The species mean average value reported by Siepmann and Finlayson, 2000 is 0.44 µg/L for diazinon acute toxicity tests accepted by CDFG. *Ceriodaphnia dubia* is the most sensitive species when the reported results for *Gammarus fasciatus* are not considered (see discussion in Section 1.2.1 below).

<sup>2</sup> The species mean average value reported by Siepmann and Finlayson, 2000 is 0.06 µg/L for chlorpyrifos acute toxicity tests. Of the freshwater species tested, *Ceriodaphnia dubia* is the most sensitive to chlorpyrifos.

**Table 4.3. Summary of potential freshwater water quality objectives derived by alternate methods**

| ALTERNATIVE                    | Diazinon                                |                   | Chlorpyrifos       |                    |
|--------------------------------|---|-------------------|--------------------|--------------------|
|                                | Acute<br>(µg/L)                         | Chronic<br>(µg/L) | Acute<br>(µg/L)    | Chronic<br>(µg/L)  |
| 1. No Change                   | 0.16 <sup>1</sup><br>0.042 <sup>2</sup> | 0.10 <sup>1</sup> | 0.025 <sup>3</sup> | 0.015 <sup>3</sup> |
| 2. No diazinon or chlorpyrifos | 0 or non detect                         | 0 or non detect   | 0 or non detect    | 0 or non detect    |
| 3. CDFG/US EPA Method          | 0.16 <sup>1</sup>                       | 0.10 <sup>1</sup> | 0.025 <sup>3</sup> | 0.015 <sup>3</sup> |

1. Regional Board staff calculations based on CDFG data set, using US EPA method. The acute criterion is a one-hour average and the chronic criterion is a four-day average – neither to be exceeded more than once every three years on the average.
2. Daily maximum based on 1/10<sup>th</sup> of the 96-hour LC50 for *Ceriodaphnia dubia*. 0.420 µg/L is found from averaging the LC50s found by CDFG (2000) and US EPA (2004).
3. CDFG (Siepmann and Finlayson, 2000) acute criterion recalculated by Regional Board staff to two significant figures per the US EPA methodology (1985).

**Table 4.4 Comparison of historical data to the alternate acute water quality objectives (1991 – 2005)**

| <b>Target</b>             | <b>1991</b>                                 | <b>1992</b>        | <b>1993</b>        | <b>1994</b>        | <b>1995</b>       | <b>1996</b>     | <b>1997</b>       | <b>1998</b>      | <b>1999</b>       | <b>2000</b>       | <b>2001</b>        | <b>2002</b>       | <b>2003</b>       | <b>2004</b>       | <b>2005</b>      |
|---------------------------|---|--------------------|--------------------|--------------------|-------------------|-----------------|-------------------|------------------|-------------------|-------------------|--------------------|-------------------|-------------------|-------------------|------------------|
| Diazinon = 0 µg/L         | <b>15%</b> <sup>a</sup><br>171 <sup>b</sup> | <b>32%</b><br>232  | <b>44%</b><br>272  | <b>54%</b><br>126  | <b>100%</b><br>14 | NS <sup>c</sup> | <b>29%</b><br>34  | <b>37%</b><br>43 | <b>25%</b><br>44  | <b>68%</b><br>95  | <b>87%</b><br>191  | <b>12%</b><br>26  | <b>1.2%</b><br>85 | <b>20%</b><br>45  | <b>86%</b><br>7  |
| Diazinon = 0.042 µg/L     | <b>9.4%</b><br>171                          | <b>20%</b><br>232  | <b>33%</b><br>272  | <b>40%</b><br>126  | <b>14%</b><br>14  | NS              | <b>8.8%</b><br>34 | <b>26%</b><br>43 | <b>9.1%</b><br>44 | <b>19%</b><br>95  | <b>17%</b><br>191  | <b>3.8%</b><br>26 | <b>0%</b><br>85   | <b>4.4%</b><br>45 | <b>0%</b><br>7   |
| Diazinon = 0.16 µg/L      | <b>0%</b><br>171                            | <b>3.0%</b><br>232 | <b>13%</b><br>272  | <b>14%</b><br>126  | <b>0%</b><br>14   | NS              | <b>0%</b><br>34   | <b>0%</b><br>43  | <b>0%</b><br>44   | <b>0%</b><br>95   | <b>5.8%</b><br>191 | <b>0%</b><br>26   | <b>0%</b><br>85   | <b>0%</b><br>45   | <b>0%</b><br>7   |
| Chlorpyrifos = 0 µg/L     | <b>1.5%</b><br>195                          | <b>11%</b><br>235  | <b>9.6%</b><br>260 | <b>23%</b><br>124  | <b>28%</b><br>14  | NS              | <b>18%</b><br>34  | <b>25%</b><br>12 | <b>41%</b><br>43  | <b>64%</b><br>96  | <b>53%</b><br>191  | <b>48%</b><br>25  | <b>17%</b><br>90  | <b>56%</b><br>45  | <b>100%</b><br>8 |
| Chlorpyrifos = 0.025 µg/L | <b>0%</b><br>195                            | <b>0.4%</b><br>235 | <b>3.5%</b><br>260 | <b>1.6%</b><br>124 | <b>7.1%</b><br>14 | NS              | <b>0%</b><br>34   | <b>0%</b><br>12  | <b>0%</b><br>43   | <b>1.0%</b><br>96 | <b>1.0%</b><br>191 | <b>16%</b><br>25  | <b>2.2%</b><br>90 | <b>2.2%</b><br>45 | <b>0%</b><br>8   |

<sup>a</sup>% of samples exceeding target<sup>b</sup>total number of samples for the year<sup>c</sup>NS = No samples analyzed during the yearData for San Joaquin River monitoring sites at:

Lander Avenue (Highway 165) near Stevinson

Hills Ferry Road near Newman

Las Palmas Avenue near Patterson

Airport Road near Vernalis

Maze Boulevard

Crows Landing

**Table 4.5. Assessment of chlorpyrifos alternatives for their consistency with Porter-Cologne and other state and federal requirements.**

| <b>Porter-Cologne Requirements</b>         | <b>No Change</b> | <b>No Chlorpyrifos</b> | <b>CDFG/ US EPA</b> |
|--|------------------|------------------------|---------------------|
| Beneficial Uses                            | +                | +                      | +                   |
| Environmental Characteristics              | 0                | 0                      | 0                   |
| Conditions Reasonably Achievable           | +                | -                      | +                   |
| Economic Considerations                    | +                | -                      | +                   |
| Need for Housing                           | 0                | 0                      | 0                   |
| Need to Recycle Water                      | 0                | 0                      | 0                   |
| <b>State and Federal Laws and Policies</b> | <b>No Change</b> | <b>No Chlorpyrifos</b> | <b>CDFG/US EPA</b>  |
| Anti-degradation                           | C                | C                      | C                   |
| Clean Water Act                            | C                | C                      | C                   |
| ESA  | C                | C                      | C                   |

Scores indicate relative degree of protection; attainability; achievability; impact or consistency with policy, as applicable, with 0 indicating neutral:

**Beneficial Uses:** Not protective of beneficial uses: - Fully protective: +

**Environmental Characteristics:** Not attainable: - Fully attainable: +

**Achievability:** Difficult to achieve - Readily achievable: +

**Economic Considerations:** Potentially significant impact: - Modest or no negative impact: +

**Housing:** Significant housing impact: - Little or no impact: +

**Recycling Water:** Significant impact on recycling water: - Little or no impact: +

C = Consistent

**Table 4.6. Assessment of diazinon alternatives for their consistency with Porter-Cologne and other state and federal requirements.**

| <b>Porter-Cologne Requirements</b>         | <b>No Change</b> | <b>No Diazinon</b> | <b>CDFG/US EPA</b> |
|--|------------------|--------------------|--------------------|
| Beneficial Uses                            | +                | +                  | +                  |
| Environmental Characteristics              | 0                | 0                  | 0                  |
| Conditions Reasonably Achievable           | +                | -                  | +                  |
| Economic Considerations                    | +                | -                  | +                  |
| Need for Housing                           | 0                | 0                  | 0                  |
| Need to Recycle Water                      | 0                | 0                  | 0                  |
| <b>State and Federal Laws and Policies</b> | <b>No Change</b> | <b>No Diazinon</b> | <b>CDFG/US EPA</b> |
| Anti-degradation                           | C                | C                  | C                  |
| Clean Water Act                            | C                | C                  | C                  |
| ESA  | C                | C                  | C                  |

Scores indicate relative degree of protection; attainability; achievability; impact or consistency with policy, as applicable, with 0 indicating neutral:

**Beneficial Uses:** Not protective of beneficial uses: - Fully protective: +

**Environmental Characteristics:** Not attainable: - Fully attainable: +

**Achievability:** Difficult to achieve: - Readily achievable: +

**Economic Considerations:** Potentially significant impact: - Modest or no negative impact: +

**Housing:** Significant housing impact: - Little or no impact: +

**Recycling Water:** Significant impact on recycling water: - Little or no impact: +

C = Consistent

**Table 4.7 River Reaches and Their Tributary Subareas**

| <b>SJR Reach</b>   | <b>Tributary Subareas</b>                    |
|--|--|
| Mendota Dam to Sack Dam  | Grassland                                    |
| Sack Dam to Lander Avenue (near Stevinson)                           | Fresno-Chowchilla, Bear Creek                |
| Lander Avenue (near Stevinson) to Hills Ferry Road (near Newman)     | Grassland, Stevinson                         |
| Hills Ferry Road (near Newman) to Las Palmas Avenue (near Patterson) | Greater Orestimba, Turlock, Merced           |
| Las Palmas Avenue (near Patterson) to Maze Boulevard                 | Westside Creeks, Northeast Bank, Tuolumne    |
| Maze Boulevard to Airport Road (near Vernalis)                       | Vernalis North, North Stanislaus, Stanislaus |



**Table 4.8 Number and Magnitude of Observed Exceedances of Proposed Loading Capacity for Combined Diazinon and Chlorpyrifos Toxicity in the mainstem SJR (2000-2004)**

| Sampling Location on the SJR     | Number of observed exceedances of proposed loading capacity for combined toxicity (2000-2004) | Average % reduction required to meet Loading Capacity for combined toxicity during observed exceedances (2000-2004) | Maximum % reduction required to meet Loading Capacity for combined toxicity during observed exceedances (2000-2004) |
|----------------------------------|---|---|---|
| near Vernalis                    | 15<br>(217 samples)   | 26%   | 40%   |
| at Maze Blvd.                    | 0<br>(20 samples)   | No observed exceedances   | No observed exceedances   |
| at Los Palmas Av. near Patterson | 1<br>(50 samples)   | 40%   | 40%   |
| at Hills Ferry Rd. near Newman   | Not Sampled   | Not Sampled   | Not Sampled   |
| near Stevinson at Lander Avenue  | 9<br>(79 samples)   | 41%   | 70%   |

**Table 4.9 Number and Magnitude of Observed Exceedances of Proposed Loading Capacity for Combined Diazinon and Chlorpyrifos Toxicity in SJR Tributaries (2000-2004)**

| Sampling Location                      | Number observed exceedances of proposed load allocations for combined toxicity / number of samples (2000-2004) | Average % reduction required to meet Loading Capacity for combined toxicity during observed exceedances (2000-2004) | Maximum % reduction required to meet Loading Capacity for combined toxicity during observed exceedances (2000-2004) |
|--|--|---|---|
| Stanislaus River at Caswell State Park | 2<br>(132 samples)   | 65%   | 76%   |
| Tuolumne River at Shiloh Road          | 9<br>(153 samples)   | 25%   | 55%   |
| Del Puerto Creek at Vineyard Road      | 18<br>(85 samples)   | 61%   | 85%   |
| Orestimba Creek at River Road          | 29<br>(155 samples)  | 76%   | 98%   |
| Merced River at River Road             | 4<br>(140 samples)   | 58%   | 70%   |
| Mud Slough near Gustine                | 2<br>(22 samples)  | 38%   | 54%   |
| Salt Slough at Lander Avenue           | 4<br>(22 samples)  | 82%   | 94%   |
| TID Lateral 5 (Harding Drain)          | 0<br>(29 samples)  | no observed exceedances   | no observed exceedances   |
| Ingram/Hospital Creeks at River Road   | 1<br>(4 samples)   | 11%   | 11%   |
| Spanish Grant Drain Near Patterson     | 0<br>(2 samples)   | no observed exceedances   | no observed exceedances   |

**Table 5.1 Summary of Differences in Dormant Season Pest Management Costs**

|                                      | <b>Almond</b>   |                               | <b>Peach</b>    |                               | <b>Apple</b>    |                               |
|--------------------------------------|-----------------|-------------------------------|-----------------|-------------------------------|-----------------|-------------------------------|
|                                      | Total cost (\$) | Percent Change from Base Case | Total cost (\$) | Percent Change from Base Case | Total cost (\$) | Percent Change from Base Case |
| <b>Base Case (diazinon)</b>          | \$2,749         | NA                            | \$3,951         | NA                            | \$11,692        | NA                            |
| <b>Base Case (chlorpyrifos)</b>      | \$2,735         | NA                            | \$3,917         | NA                            | \$11,688        | NA                            |
| <b>Scenario 1</b>                    | \$2,750         | 1%                            | \$3,937         | 0%                            | \$11,673        | 0%                            |
| <b>Scenario 2</b>                    | \$2,778         | 2%                            | \$4,000         | 1%                            | \$11,741        | 0%                            |
| <b>Scenario 3</b>                    | \$2,760         | 1%                            | \$3,962         | 1%                            | \$11,703        | 0%                            |
| <b>Scenario 4</b>                    | \$2,898-\$2,909 | 6%                            | \$4,078         | 3%                            | \$11,832        | 1%                            |
| <b>Percent change from Base Case</b> |                 | 1% to 6%                      |                 | 0% to 3%                      |                 | 0% to 1%                      |

**Table 5.2 Surface Irrigation - Initial Capital Cost and Recurring Maintenance Costs**  
(from Burt et al. 2000)

| <b>System Type</b> | <b>Capital \$/acre</b> | <b>Maintenance \$/acre/year</b> | <b>Labor hrs/acre</b> | <b>Energy kwh/ac-in</b> |
|--------------------|------------------------|---------------------------------|-----------------------|-------------------------|
| Basin Irrigation   | 3192                   | 51                              | 0.3                   | n/a                     |
| Border Strip       | 2228                   | 51                              | 0.4                   | n/a                     |
| Contour Ditch      | 140                    | 13                              | 2.5                   | n/a                     |
| Continuous Flood   | 1010                   | 26                              | 0.3                   | n/a                     |
| Furrow             | 1703                   | 51                              | 1                     | n/a                     |
| Corrugation        | 1475                   | 51                              | 1.25                  | n/a                     |

**Table 5.3 Sprinkler Irrigation - Initial Capital Cost and Recurring Maintenance Costs**  
(from Burt et al. (2000))

| <b>System Type</b>    | <b>Capital<br/>\$/acre</b> | <b>Maintenance<br/>\$/acre/year</b> | <b>Labor<br/>hrs/acre</b> | <b>Energy<br/>kwh/ac-in</b> |
|-----------------------|----------------------------|-------------------------------------|---------------------------|-----------------------------|
| Hand Move Lateral     | 225                        | 5                                   | 0.175                     | 15.4                        |
| End Row Lateral       | 325                        | 10                                  | 0.103                     | 15.4                        |
| Side Roll Lateral     | 388                        | 8                                   | 0.123                     | 15.4                        |
| Traveling Gun         | 450                        | 27                                  | 0.072                     | 43.2                        |
| Center Pivot          | 363                        | 18                                  | 0.01                      | 16.5                        |
| Center Pivot w/corner | 450                        | 27                                  | 0.01                      | 17.5                        |
| Linear Move w/ditch   | 488                        | 29                                  | 0.021                     | 16.5                        |
| Linear Move w/pipe    | 738                        | 44                                  | 0.021                     | 19.5                        |
| Portable Solid Set    | 1200                       | 24                                  | 0.103                     | 15.4                        |
| Permanent Solid Set   | 1163                       | 12                                  | 0.01                      | 15.4                        |

156

**Table 5.4 Micro-irrigation - Initial Capital and Maintenance Costs**  
(from Burt et al (2000))

| <b>System Type</b>       | <b>Capital<br/>\$/acre</b> | <b>Maintenance<br/>\$/acre/year</b> | <b>Labor<br/>hrs/acre</b> | <b>Energy<br/>kwh/ac-in</b> |
|--------------------------|----------------------------|-------------------------------------|---------------------------|-----------------------------|
| Drip Vineyards           | 1050                       | 105                                 | 0.04                      | 10.95                       |
| Drip Orchards Surface    | 850                        | 85                                  | 0.04                      | 10.95                       |
| Drip Orchards Subsurface | 1100                       | 110                                 | 0.04                      | 10.95                       |
| Micro Orchards           | 950                        | 95                                  | 0.04                      | 10.95                       |
| Drip Row Surface         | 700                        | 70                                  | 0.04                      | 10.95                       |
| Drip Row Subsurface      | 1700                       | 170                                 | 0.04                      | 10.95                       |

**Table 5.5. Estimated Cost to Convert from Flood to Sprinkler Irrigation**

|  | <b>Almonds</b>       |                      | <b>Walnuts</b>      |                     |
|--|----------------------|----------------------|---------------------|---------------------|
| Total acres in TMDL area   | 231,788 <sup>1</sup> | 231,788 <sup>1</sup> | 28,057 <sup>1</sup> | 28,057 <sup>1</sup> |
| Estimated percent in flood irrigation  | 40% <sup>2</sup>     | 60% <sup>3</sup>     | 40% <sup>2</sup>    | 60% <sup>3</sup>    |
| Estimated acres in flood irrigation  | 92,715               | 139,073              | 11,223              | 16,834              |
| Estimated percentage acres using diazinon or chlorpyrifos                                    | 30% <sup>1</sup>     | 30% <sup>1</sup>     | 65% <sup>1</sup>    | 65% <sup>1</sup>    |
| Increased cost/acre for sprinkler irrigation   | \$196                | \$196                | \$196               | \$196               |
| Total increased cost to convert acres using diazinon or chlorpyrifos to sprinkler irrigation | \$5,451,654          | \$8,177,492          | \$1,429,810         | \$2,144.652         |

1.From 2002 PUR

2 Estimate based on information from USDA 1998 Farm and Ranch Irrigation Survey

3 Estimate based on information from Zoldoske. 2002

**Table 5.6 Drainage Practices Capital and Maintenance Costs**

| <b>System Type</b>             | <b>Capital<br/>\$/acre</b> | <b>Maintenance<br/>\$/acre/year</b> |
|--------------------------------|----------------------------|-------------------------------------|
| Surface Drainage Recirculation | 812                        | 55                                  |
| Temporary Retention Ponds      | 340                        | 50                                  |

**Table 5.7 Summary of Differences in Irrigation Season Pest Management Costs**

|                                      | <b>Almond</b>   |                               | <b>Alfalfa</b>  |                               |
|--------------------------------------|-----------------|-------------------------------|-----------------|-------------------------------|
|                                      | Total cost (\$) | Percent Change from Base Case | Total cost (\$) | Percent Change from Base Case |
| <b>Base Case (chlorpyrifos)</b>      | \$2,781         | NA                            | \$1,009         | NA                            |
| <b>Scenario 1</b>                    | \$2,884         | 3.7%                          | \$1,124         | 11.4%                         |
| <b>Scenario 2</b>                    | \$2,871         | 3.2%                          | \$1,084         | 7.4%                          |
| <b>Scenario 3</b>                    | \$2,899         | 4.2%                          |                 |                               |
| <b>Percent change from Base Case</b> |                 | 3.2% to 4.2%                  |                 | 7.4% to 11.4%                 |

158

**Table 7.1 Summary of Public Workshops**

| <b>Date</b>    | <b>Workshop</b>   |
|----------------|---|
| August 2000    | Initial Outreach of OP Pesticide TMDL                           |
| November 2000  | Initial Stage of the TMDL Development / Draft Problem Statement |
| January 2001   | Introduced Elements of TMDL and Monitoring Data                 |
| June 2001      | Draft Numeric Target Report                                     |
| March 2002     | Draft Source Analysis Report                                    |
| July 2002      | Draft TMDL Report   |
| September 2002 | Draft TMDL Implementation Framework                             |
| January 2005   | CEQA Scoping Meeting and Public Workshop                        |